



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

PROBLEMS.

7. Proposed by Professor J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

To determine the function $F(x)$ so that $F(x+y) \times F(x-y) = [F(x)]^2 - [F(y)]^2$.

8. A woodman fells a tree 2 feet in diameter, cutting half way through from each side. The lower face of each cut is horizontal, and the upper face makes an angle of 45° with the horizontal. How much wood does he cut out?

[Selected from *Byerly's Integral Calculus*.]

9. Proposed by Professor G. B. M. ZERR, A. M., Principal of Schools, Staunton, Virginia.

The solids bounded by the surfaces whose equations are $\left|\frac{x}{a}\right|^{\frac{2}{3}} + \left|\frac{y}{b}\right|^{\frac{2}{3}} + \left|\frac{z}{c}\right|^{\frac{2}{3}} = 1$, and $x^{\frac{2}{3}} + y^{\frac{2}{3}} + z^{\frac{2}{3}} = b^{\frac{2}{3}}$ where $a > b > c$ have their centers coincident. Find (1 and 2) the volume of each without the other, and (3) the volume common to both by direct integration, using the formula $V = \iiint dx dy dz$.

- 10 Proposed by ERIC DOOLITTLE, Instructor in Mathematics, State University of Iowa.

Prove or disprove the following theorem: If O be any circle, and AB any straight line either within or without the circumference, and if a perpendicular be dropped from O upon AB and prolonged backward to meet the circumference in P , then will the angle whose vertex lies at P and whose sides pass through A and B , cut a portion CPD from the circle which shall be greater than that cut by any other angle whose vertex lies on the circumference and whose sides pass through A and B .

[If any one can give a solution without the use of the Calculus, it will also be acceptable —ED.]

11. Proposed by H. W. DRAUGHON, Clinton, Louisiana.

A ribbon, 1 inch wide is wrapped spirally around a right prism, altitude 10 ft., bases of n sides, radius of circumscribed circle, 1 ft., so as to cover the entire convex surface. (1) What is the length of the ribbon? (2) If the ribbon is unwound and kept tense, by a power acting on the lower end, and moving in the plane of the lower base, what will be the length of the curve described by the power?

MECHANICS.

Conducted by B. F. FINKEL, Kidder, Missouri. All contributions to this department should be sent to him.

PROBLEMS.

6. Proposed by THOMAS W. WRIGHT, M. A., Ph. D., Professor of Applied Mathematics and Physics, Union College, Schenectady, New York.